

Mammary Tumors in Dogs - Don't Ignore a Lump!

Each October in the United States, various organizations strive to raise awareness of and funds for finding a cure for breast cancer in humans. Breast cancer, also called mammary cancer, in dogs has both similarities and differences to the human condition. To end the month of October, I review the diagnosis and treatment of mammary tumors in dogs. Happy reading!

Mammary Tumors: Basic Anatomy & Risk Factors

Just as humans are encouraged to perform self-examinations of breast tissue in an effort to identify tumors when they're small, so too should pet parents frequently evaluate their pets' breast tissue. This requires a basic knowledge of a dog's mammary tissue. Female dogs have 5 pairs of mammary glands

- 2 cranial thoracic glands
- 2 caudal thoracic glands
- 2 cranial abdominal glands
- 2 caudal abdominal glands
- 2 inguinal glands




Illustration of the mammary

glands in dogs. Image courtesy
of www.marvistavet.com

Approximately 60-70% of mammary tumors in dogs in either the caudal abdominal or inguinal mammary glands. Mammary tumors are the most common tumor seen in female dogs, and the risk for developing a breast tumor increases with age in those who have not undergone ovariohysterectomy (been spayed):

- 0.05% risk if spayed before 1st heat
- 8% risk if spayed after 1st heat
- 26% risk if spayed after 2nd heat

Multiple studies suggest hormones like estrogen and progesterone likely play a role in the development of mammary gland tumors. Thus the use of hormones therapeutically also increases the risk of benign mammary tumors in dogs.

Mammary Tumors: Getting A Diagnosis

Approximately 50% of all breast tumors are cancerous, and about 50% of them will spread or metastasize. Remember our [previous discussion](#) about when pet parents should have a skin mass evaluated by a family veterinarian:

- When the mass has been present for more than 1 month
- When the mass is larger than the size of a pea

Many masses associated with or under the skin (also called subcutaneous) are initially evaluated with a simple and non-invasive procedure called a fine needle aspiration or FNA. Unfortunately, FNAs of mammary tumors do not reliably differentiate between non-cancerous and cancerous disease. Nevertheless, an FNA is still quite helpful, as it can rule out other types of skin and subcutaneous masses.



A large mammary tumor in a dog

If the results of an FNA test identify mammary tissue, then the mass should be surgically removed to determine definitively if it is cancerous or benign.

Surgery is the initial gold standard therapy for dogs with mammary tumors.

Depending on the size of the breast tumor, the surgical procedure can be minor or extensive. Possible surgeries include:

- Lumpectomy: removal of small nodules and associated skin
- Mastectomy: removal of a single mammary gland
- Regional mastectomy: removal of two or more adjacent mammary glands
- Unilateral mastectomy: removal of all mammary glands on one side
- Bilateral mastectomy: removal of all mammary glands - each side is removed ~4 weeks apart
- Radical mastectomy: removal of a large mammary tumor and extensive tissue around the mass because the tumor has adhered to the abdominal wall

The regional lymph nodes (axillary and/or inguinal lymph nodes) are also often removed, especially if they are enlarged or firm, to screen for evidence of metastasis. Consultation with a board-certified veterinary surgeon can be invaluable for determining the best surgical procedure for your fur baby.

Performing an ovariohysterectomy (spaying) at the time of mammary tumor removal is currently quite controversial. Multiple studies have shown spaying does not affect prognosis. More recently, however, two studies ([study #1](#); [study #2](#)) documented increased survival in dogs spayed at the time their mammary tumors were removed or less than two years of the time of their mammary tumor removal.

Mammary Tumors: Staging

Staging is simply the medical term used to describe a series of tests performed to determine how much the body is affected by cancer. The staging tests for mammary cancer are non-invasive, and include:

- Complete physical examination
- Blood and urine evaluations
- Chest radiographs/x-rays
- Abdominal sonography

Some of these staging tests are performed prior to surgery. For example, simple blood and urine tests are recommended [prior to any anesthesia and surgery](#). Additionally many veterinarians obtain chest radiographs/x-rays prior to surgery to empower pet parents with information about the presence of lung metastases.

Mammary Tumors: Treatment After Surgery

Following surgery, additional treatment, including chemotherapy, may be recommended. Recall our previous discussion about [chemotherapy in dogs and cats](#). Simply stated anti-cancer drug therapy is a much friendlier process in veterinary medicine than it is for humans. Many factors influence the decision to pursue chemotherapy, including:

- Presence of metastasis
- Specific changes noted in the biopsy
- Tumor grade (a way describing the biologic aggressiveness of a cancer)
- Prior et parent experience with chemotherapy

There are many chemotherapy protocols that have been investigated in dogs with mammary cancer. Recently a drug called paclitaxel ([Paccal Vet®](#)) received approval from the US Food and Drug Administration for treatment of non-

resectable stage III, IV and V mammary carcinoma in dogs. Unquestionably consultation with a board-certified veterinary oncologist (or a board-certified veterinary internal medicine specialist if an oncologist is not available) is strongly recommended to empower families with the most up-to-date information they need to make an informed decision about chemotherapy.

Tumor stage has also been shown to affect survival time. For example, dogs with stage I disease have reported 97.9% 1-year and 2-year survival times while those with stage IV disease have 13.6% 1-year and 2-year survival times. Tumor size also seems to affect prognosis. Those dogs with breast tumors less than three centimeters in diameter have significantly better prognoses. In another study, patients with tumors larger than five centimeters in diameter were more likely to develop metastases than those with smaller masses; these patients were also seven times more likely to die within seven years. Furthermore, studies have shown dogs with no metastases were more than three times more likely to be alive one year after diagnosis.

Several other factors have been associated with a poor prognosis for dogs with mammary cancer, particularly:

- Certain types of mammary cancers (inflammatory carcinoma, ductular carcinoma, carcinosarcoma)
- Degree of ulceration of the mammary tumor
- High grade
- Invasion of the cancer into blood vessels or lymph vessels within the mammary tissue
- Duration of tumor presence

The take-away message about mammary tumors in dogs...

Mammary or breast tumors are extremely common tumors in female dogs. Early identification and removal are essential, as surgery can be curative in ~75% of affected patients. Consulting and partnering with board-certified veterinary specialists in surgery and cancer treatment can be uniquely helpful for developing efficient and effective diagnostic and treatment plans.

To find a board-certified veterinary surgeon, please visit the [American College of](#)

[Veterinary Surgeons.](#)

To find a board-certified veterinary oncologist, please visit the [American College of Veterinary Internal Medicine.](#)

Wishing you wet-nosed kisses,

cgb